



INDIVIDUALIZED SOLUTIONS FOR INTEGRATIVE CLINICIANS™

[FAQ](#) | [Ordering Info](#) | [Site Map](#) | [Contact us](#)

[Print](#)

[Home](#)

[Clinicians](#)

[Patients / Consumers](#)

[International](#)

[Test Menu](#)

[Introduction](#)

[Table of Contents](#)

[View by Category](#)

[CPT Codes](#)

[How to Order](#)

[Learning Center](#)

[News and Events](#)

[About Metamatrix](#)

[Contact Us](#)

[Web Store](#)

[Search](#)

Clinician login:

Email

Password

## Detoxification Capacity - Saliva & Urine

[Overview](#)

[Clinician Info](#)

[CPT Codes](#)

[Kit Instructions](#)

[Sample Reports](#)

[Interpretive Guide](#)

[References](#)



Three compounds, caffeine, acetaminophen, and acetylsalicylic acid, are taken orally to challenge the liver's Phase I and Phase II detoxification capacity. Saliva and urine are analyzed for metabolites of the three compounds to determine how well the liver can convert and clear toxins from the body. This test is not recommended for children 12 years old and younger.

***\*Profile is not available in New York***



Find at Metamatrix.com:

[Cardiovascular health](#)

[GI Effects](#)

[Organic acids](#)

[Celiac disease](#)

[Amino acid](#)

[Vitamin D](#)

[IgG antibodies](#)

[Bone resorption](#)

[Breast cancer](#)

[Yeast](#)

[Estrogen metabolites](#)

[Homocysteine](#)

[Autism](#)

[Oxidative stress](#)

[Case studies](#)

© 2008 Metamatrix, Inc. All rights reserved.